

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C11D 1/94, 3/37, A61K 7/50, 7/48	A1	(11) International Publication Number: WO 97/40125 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/EP97/01147 (22) International Filing Date: 6 March 1997 (06.03.97) (30) Priority Data: 08/637,144 24 April 1996 (24.04.96) US (71) Applicant (for AU BB CA GB IE IL KE LC LK LS MN MW NZ SD SG SZ TT UG only): UNILEVER PLC (GB/GB); Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AU BB CA GB IE IL KE LC LK LS MN MW NZ SD SG SZ TT UG): UNILEVER N.V. (NL/NL); Weena 455, NL-3013 AL Rotterdam (NL). (72) Inventors: HE, Mengtao; 9 Tuxedo Drive, Wayne, NJ 07470 (US). FAIR, Michael, Joseph; 290 Anderson, Hackensack, NJ 07601 (US). MASSARO, Michael; 39 Dover Road, Congers, NY 10920 (US). (74) Agent: MOLE, Peter, Geoffrey; Unilever plc, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: LIQUID COMPOSITIONS COMPRISING HYDROPHOBICALLY MODIFIED POLYALKYLENE GLYCOLS AS MILD- NESS ACTIVES (57) Abstract <p>The present invention relates to liquid detergent compositions comprising anionic/amphoteric surfactant systems. Low addition levels of specific hydrophobically modified polyalkylene glycol wherein ratio of hydrophobically modified polyalkylene glycol to anionic surfactant is defined have been found to remarkably enhance mildness without sacrificing processibility and desired user properties. In a second embodiment, the invention relates to a method for enhancing mildness in liquid detergent compositions comprising anionic surfactant by adding said defined hydrophobically modified polyalkylene glycols.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

LIQUID COMPOSITIONS COMPRISING
HYDROPHOBICALLY MODIFIED POLYALKYLENE GLYCOLS
AS MILDNESS ACTIVES

5

FIELD OF THE INVENTION

The present invention relates to liquid personal wash compositions (e.g., shower gels), particularly, to compositions comprising (1) one or more anionic surfactants and (2) one or more amphoteric surfactants. Specifically, the invention relates to the incorporation of relatively low addition levels of specific hydrophobically modified polyalkylene glycols (HMPEG) in the liquids. Through careful balancing of anionic surfactant to HMPEG and specific selection of the HMPEG, enhanced mildness is obtained without sacrificing processability, lather performance and other desired user properties.

20

BACKGROUND

The use of hydrophobically modified polyalkylene glycols in liquid personal wash compositions is not new. High molecular weight hydrophobically modified polyalkylene glycols (HMPEG) have been used previously as thickeners in liquids at an addition level of less than 5% wt. of total composition; beyond this level, over-thickening can be induced. Some prior art (e.g., Patents by Egan et al. discussed below) teach that only relatively high levels of addition of HMPEG (HMPEG to anionic surfactant weight ratio between 1:1 and 4:1) can effectively reduce the skin irritation potential of anionic surfactants. However, this high levels of HMPEG can cause over-thickening, and the referred patents also teach the solution to control the viscosity. Two GB patent application by Dias et al. teach that the combination of urea, alpha- or beta-hydroxy acid,

and animal/oil derived ethoxylated nonionic surfactants (some of them are the HMPEG of the subject invention) are needed to achieve desired skin feel effect.

5 The references are discussed below.

Patent No. WO 9,423,695 to Unilever NV teaches a personal washing formulation that contains 3-15% sodium lauryl (EO) sulfate, 3-10% sulphosuccinate surfactants, 0.5-3% thickener such as fatty acid glyceride polyglycol ether (a HMPEG), and 0.1-5% of an emulsifier such as PEG 40 hydrogenated castor oil. The percentage of the fatty acid glyceride polyglycol ether is outside the range of choice of our invention, and the molecular weight of PEG 40 hydrogenated castor oil is below 4000, which is also outside the range of choice of our invention.

Patent No. SU 1,330,148 to S. Navardausk, R. Zilberman, and D. Kurdyumova teaches a liquid cleanser formulation containing 2-4 % wt. of polyethylene glycol ester of 10-16C (a HMPEG) as thickener mixed with other ingredients. The percentage of the 10-16C polyethylene glycol ester in total composition is below 5%, which is outside the range of choice of our invention.

Patent No. DE 4,409,189 to Chem-Y Chem Fab GMBH teaches a surfactant composition containing 5-20% anionic surfactants, 2.5-10% alkyl polyglucoside, 1-4% ethoxylated rape seed oil fatty acid (can be a HMPEG), and 0.5-2.5% polyoxyethylene propylene dioleate (a HMPEG) as thickener. The percentage of the HMPEG in total composition is outside the range of choice of our invention.

Patent No. FR 2,336,475 to Gillette Co. teaches the use of amphoteric surfactant and nonionic surfactant as major

actives and anionic surfactant as coactives in a PW liquid. The formulation contains 2% thickener such as polyethylene glycol 6000 distearate (a HMPEG). The percentage of the HMPEG in total composition is outside the range of choice of our invention.

Patent No. US 4,828,750 to F. Simion, L. Rhein, J. Blake-Haskins, S. Babulak and R. Cantore teaches a formulation containing 0.25-6% nonionic surfactant, 0.05-5% of an organic acid, 0.1-4% of a thickening agent, such as a diester of stearic acid and polyoxyethylene (can be a HMPEG of the subject invention). The level of the diester of stearic acid and polyoxyethylene is below 5% by wt. total composition, which is outside the range of choice of our invention.

US Patent No.s 4,247,425, 4,343,726, and 4,256,611 to R. Egan teach liquid skin cleansing formulations containing anionic surfactant and hydrophobically modified polyalkylene glycols (e.g., POE(200)-glyceryl-stearate) as mildness enhancers. These patents showed that only at relatively high addition level of the hydrophobically modified polyalkylene glycols (hydrophobically modified polyalkylene glycols / anionic surfactant weight ratio above 1:1 (preferably 1:1 to 4:1)), the hydrophobically modified polyalkylene glycols can significantly reduce the irritation of anionic surfactant. However, use of high levels of HMPEG may thicken the liquid composition and require extra efforts to control the viscosity of the formulation, which is part of the art of this invention.

UK Patent Application Nos. GB 2,288,811A and GB 2, 288, 812A to L. Dias, M. Giret, C. Leahy, and R. Elliot teach liquid skin cleansing formulations containing 5 to 50% of anionic, amphoteric and zwitterionic surfactants in general,

0.1 to 20% of a soluble or dispersible nonionic surfactant selected from ethoxylated animal and vegetable oils and fats and mixtures thereof, 0.5-8% urea, and 0.1-5% alpha or beta hydroxy acid. Evidently, the art of this invention is to use
5 a combination of urea, the animal/vegetable oil derived ethoxylated nonionic surfactants (some of them can be the HMPEG of the subject invention), and hydroxy acid to achieve desirable skin feel and mildness (PG11, Line 4 to 6, and PG4, Line 26 to 28). However, neither in-vivo nor in-vitro data
10 were presented to support the mildness claim, and no anionic/ethoxylated nonionic weight ratio was specified in these applications. Finally, the references require a specific combination of ingredients (e.g., urea and the oil-fat derived ethoxylated nonionic surfactants) to achieve the
15 desired skin feel effects, thus use of urea is a criticality. In contrast, the use of urea is not required in the subject invention, and, in fact, is specifically disclaimed.

The subject invention differs from the prior art
20 referred above, alone or in combination, in that the applicants have found that at relatively low levels, defined hydrophobically modified polyalkylene glycols (e.g., having specific HMPEG to anionic surfactant weight ratio below 1:1, specific range of molecular weight, and specific ethylene
25 oxide : hydrophobe ratio) in specific surfactant systems (e.g., containing greater than or equal to 50% anionic surfactant(s); and also necessarily containing at least some amphoteric surfactant) will result in enhanced mildness of the specific systems without sacrificing processibility and
30 user properties. Additionally, the low levels of addition of HMPEG avoids the problem of over-thickening and can reduce the cost of the liquid compositions.

BRIEF SUMMARY OF THE INVENTION

Unexpectedly, applicants have found that in liquid personal wash compositions comprising a surfactant system comprising:

- (1) 3% to 30% by wt., preferably 5% to 20%, total composition anionic surfactant or mixture of anionic surfactants, wherein anionic is greater than or equal to 50% of the surfactant system; and
- (2) 0.1 to 20% by wt. total composition one or more amphoteric surfactants;
- (3) the addition of 5 to 25% by wt. (preferably 5 to 15%) of total composition of hydrophobically modified polyalkylene glycols (HMPEG) wherein weight of alkylene oxide (e.g., ethylene oxide) portion is between 60% and 90%, preferably 85% to 97% wt. of the weight of the HMPEG, will lead to significantly enhanced mildness in such compositions without sacrificing processibility and desired user properties, such as rich and creamy lather.

Weight ratio of HMPEG to anionic surfactant is between 1:1.1 to 1:10. This ratio is a criticality because above the range of this ratio, special efforts have to be made to control the high viscosity of liquid personal wash compositions, and the cost of the composition can be raised; below this range, skin irritation potential of anionic surfactant can not be effectively mitigated.

The HMPEG should have molecular weight of 4,000 to 25,000, preferably 4,000 to 15,000. This MW range is a criticality because, above this range of molecular weight, HMPEG can cause over-thickening and cause processing and handling difficulties. Below this MW range, HMPEG approaches the structure of a conventional ethoxylated nonionic

detergent and may impair lather performance and other desired user properties.

BRIEF DESCRIPTION OF THE DRAWINGS

5

Figure 1a and Figure 1b show the Zein % dissolved by acyl isethionate / cocoamido propyl betaine as a function of nonionic polymeric surfactant concentration. In contrast to PEG 8000, POE(200) glyceryl stearate and POE(200) glyceryl
10 tallowate significantly reduced the Zein % dissolved at relatively low levels. Therefore the irritation potential of a personal washing bar can be further reduced by including relatively low levels (i.e. less than 30%) of defined hydrophobically modified polyalkylene glycols in a full
15 liquid composition.

Figure 2 shows that, even at relatively low levels of addition, a HMPEG of the invention (POE(200) glyceryl stearate) significantly reduces skin irritation caused by
20 sodium acyl isethionate.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to novel liquid personal
25 water compositions, particularly compositions in which the surfactant system comprises greater than 50% of the surfactant system anionic surfactant or surfactants, and additionally comprises one or more amphoteric surfactants.

30 Unexpectedly, applicants have found that when relatively small amounts (5 to 25% by wt. composition) of a defined hydrophobically modified polyalkylene glycols (i.e., defined by MW of 4,000-25,000; by percentage of EO of the polymer being between 60% and 99% wt.;) is used, and when there is a
35 defined ratio of the hydrophobically modified polymer to

anionic surfactant (i.e., 1:1.1 to 1:10), the liquid composition is significantly milder (as defined by zein dissolution and patch tests) than either in the absence of the polymer or if a different alkylene oxide (e.g., polyethylene oxide) is used.

The compositions are defined in greater detail below:

Surfactant System

The surfactant system of the subject invention generally comprises 5 to 50% by weight, preferably 10 to 40% by wt. of the composition and comprises:

- (a) 3% to 30%, preferably 5 to 20% by wt. total composition one or more anionic surfactants wherein the anionic surfactant comprises 50% or more of the surfactant system;
- (b) 0.1 to 20% by wt., preferably 3% to 10% total composition amphoteric and/or zwitterionic surfactant; and
- (c) 0% to 10% optional nonionic surfactant (other than hydrophobically modified polyalkylene glycols of invention).

The anionic surfactant may be, for example, an aliphatic sulfonate, such as a primary alkane (e.g., C_8-C_{22}) sulfonate, primary alkane (e.g., C_8-C_{22}) disulfonate, C_8-C_{22} alkene sulfonate, C_8-C_{22} hydroxyalkane sulfonate or alkyl glyceryl ether sulfonate (AGS); or an aromatic sulfonate such as alkyl benzene sulfonate.


The anionic may also be an alkyl sulfate (e.g., $C_{12}-C_{18}$ alkyl sulfate) or alkyl ether sulfate (including alkyl glyceryl ether sulfates). Among the alkyl ether sulfates are those having the formula:



wherein R is an alkyl or alkenyl having 8 to 18 carbons, preferably 12 to 18 carbons, n has an average value of greater than 1.0, preferably between 2 and 3; and M is a solubilizing cation such as sodium, potassium, ammonium or substituted ammonium. Ammonium and sodium lauryl ether sulfates are preferred.

The anionic may also be alkyl sulfosuccinates (including mono- and dialkyl, e.g., C₈-C₂₂ sulfosuccinates); alkyl and acyl taurates, alkyl and acyl sarcosinates, sulfoacetates, C₈-C₂₂ alkyl phosphates and phosphates, alkyl phosphate esters and alkoxyl alkyl phosphate esters, acyl lactates, C₈-C₂₂ monoalkyl succinates and maleates, sulphoacetates, and acyl isethionates.

Sulfosuccinates may be monoalkyl sulfosuccinates having the formula:




amido-MEA sulfosuccinates of the formula



wherein R⁴ ranges from C₈-C₂₂ alkyl and M is a solubilizing cation;


amido-MIPA sulfosuccinates of formula



where M is as defined above.

Also included are the alkoxyated citrate
sulfosuccinates; and
alkoxyated sulfosuccinates such as the following:


5



wherein $n = 1$ to 20; and M is as defined above.

10


Sarcosinates are generally indicated by the formula



wherein R ranges from C_8 - C_{20} alkyl and M is a
solubilizing cation.

15

Taurates are generally identified by formula




20

wherein R^2 ranges from C_8 - C_{20} alkyl, R^3 ranges from C_1 - C_4
alkyl and M is a solubilizing cation.

Another class of anionics are carboxylates such as
follows:

25



wherein R is C_8 to C_{20} alkyl; n is 0 to 20; and M is as
defined above.

30


Another carboxylate which can be used is amido alkyl
polypeptide carboxylates such as, for example, Monteine LCQ^(R)
by Seppic.

35

Another surfactant which may be used are the C₆-C₁₈ acyl isethionates. These esters are prepared by reaction between alkali metal isethionate with mixed aliphatic fatty acids having from 6 to 18 carbon atoms and an iodine value of less than 20. At least 75% of the mixed fatty acids have from 12 to 18 carbon atoms and up to 25% have from 6 to 10 carbon atoms.

Acyl isethionates, when present, will generally range from about 0.5-15% by weight of the total composition. Preferably, this component is present from about 1 to about 10%.

The acyl isethionate may be an alkoxyated isethionate such as is described in Ilardi et al., U.S. Patent No. 5,393,466, hereby incorporated by reference into the subject application. This compound has the general formula:




wherein R is an alkyl group having 8 to 18 carbons, m is an integer from 1 to 4, X and Y are hydrogen or an alkyl group having 1 to 4 carbons and M' is a monovalent cation such as, for example, sodium, potassium or ammonium.

In general the anionic component will comprise from about 1 to 20% by weight of the composition, preferably 5 to 15%, most preferably 5 to 12% by weight of the composition.

Zwitterionic and Amphoteric Surfactants

Zwitterionic surfactants are exemplified by those which can be broadly described as derivatives of aliphatic

quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight or branched chain, and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. A general formula for these compounds is:



wherein R^2 contains an alkyl, alkenyl, or hydroxy alkyl radical of from about 8 to about 18 carbon atoms, from 0 to about 10 ethylene oxide moieties and from 0 to about 1 glyceryl moiety; Y is selected from the group consisting of nitrogen, phosphorus, and sulfur atoms; R^3 is an alkyl or monohydroxyalkyl group containing about 1 to about 3 carbon atoms; X is 1 when Y is a sulfur atom, and 2 when Y is a nitrogen or phosphorus atom; R^4 is an alkylene or hydroxyalkylene of from about 1 to about 4 carbon atoms and Z is a radical selected from the group consisting of carboxylate, sulfonate, sulfate, phosphonate, and phosphate groups.

25

Examples of such surfactants include:

4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate;

30 5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxypentane-1-sulfate;

3-[P,P-diethyl-P-3,6,9-trioxatetradecylphosphonio]-2-hydroxypropane-1-phosphate;

3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropylammonio]-propane-1-phosphonate;

35 3-(N,N-dimethyl-N-hexadecylammonio)propane-1-sulfonate;

3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxypropane-1-

sulfonate;

4-[N,N-di(2-hydroxyethyl)-N-(2-hydroxydodecyl)ammonio]-
butane-1-carboxylate;


3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-
propane-1-phosphate;

3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-
phosphonate; and

5-[N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-
hydroxy-pentane-1-sulfate.

10

Amphoteric detergents which may be used in this
invention include at least one acid group. This may be a
carboxylic or a sulphonic acid group. They include
quaternary nitrogen and therefore are quaternary amido acids.
They should generally include an alkyl or alkenyl group of 7
to 18 carbon atoms. They will usually comply with an overall
structural formula:



25

where R^1 is alkyl or alkenyl of 7 to 18 carbon atoms;

R^2 and R^3 are each independently alkyl, hydroxyalkyl or
carboxyalkyl of 1 to 3 carbon atoms;

30

n is 2 to 4;

m is 0 to 1;


35

X is alkylene of 1 to 3 carbon atoms optionally


substituted with hydroxyl, and

Y is $-\text{CO}_2-$ or $-\text{SO}_3-$

5 Suitable amphoteric detergents within the above general formula include simple betaines of formula:




and amido betaines of formula:



where m is 2 or 3.


In both formulae R^1 , R^2 and R^3 are as defined previously. R^1 may in particular be a mixture of C_{12} and C_{14} alkyl groups derived from coconut so that at least half, preferably at least three quarters of the groups R^1 have 10 to 14 carbon atoms. R^2 and R^3 are preferably methyl.

A further possibility is that the amphoteric detergent is a sulphobetaine of formula




14

or




where m is 2 or 3, or variants of these in which -
 $(CH_2)_3 SO_3^-$ is replaced by




In these formulae R^1 , R^2 and R^3 are as discussed
 previously.


A further possibility is that the amphoteric detergent
 is a sulphobetaine of formula



or



where m is 2 or 3, or variants of these in which -
 $(CH_2)_3 SO_3^-$ is replaced by



5 In these formulae R^1 , R^2 and R^3 are as discussed previously.

10 Amphoacetates and diamphoacetates are also intended to be covered in possible zwitterionic and/or amphoteric compounds which may be used.

15 The amphoteric/zwitterionic generally comprises 0.1 to 20% by weight, preferably 0.1% to 15%, more preferably 0.1 to 10% by wt. of the composition.


 In addition to one or more anionic and amphoteric and/or zwitterionic, the surfactant system may optionally comprise a nonionic surfactant.

20 The nonionic which may be used includes in particular the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone or with propylene
25 oxide. Specific nonionic detergent compounds are alkyl (C_6 - C_{22}) phenols-ethylene oxide condensates, the condensation products of aliphatic (C_8 - C_{18}) primary or secondary linear or branched alcohols with ethylene oxide, and products made by
30 condensation of ethylene oxide with the reaction products of propylene oxide and ethylenediamine. Other so-called nonionic detergent compounds include long chain tertiary amine oxides, long chain tertiary phosphine oxides and dialkyl sulphoxides.

The nonionic may also be a sugar amide, such as a polysaccharide amide. Specifically, the surfactant may be one of the lactobionamides described in U.S. Patent No. 5,389,279 to Au et al. which is hereby incorporated by reference or it may be one of the sugar amides described in Patent No. 5,009,814 to Kelkenberg, hereby incorporated into the subject application by reference.

Other surfactants which may be used are described in U.S. Patent No. 3,723,325 to Parran Jr. and alkyl polysaccharide nonionic surfactants as disclosed in U.S. Patent No. 4,565,647 to Llenado, both of which are also incorporated into the subject application by reference.

Preferred alkyl polysaccharides are alkylpolyglycosides of the formula



wherein R^2 is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 0 to 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from 1.3 to about 10, preferably from 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4- and/or 6-position, preferably predominantly the 2-position.

Nonionic comprises 0 to 10% by wt. of the composition.

Hydrophobically Modified Polyalkylene Glycols (HMPEG)

5 The hydrophobically modified polyalkylene glycols
(HMPEG) of the subject invention are generally commercially
available nonionic polymeric surfactants having a broad
molecular weight range from about 4000 to 25000 (preferably
4000 to 15000). This range of molecular weight is a
10 criticality because, above the defined range of molecular
weight, HMPEG can make liquid formulations sticky and
viscous, which causes processing problems, such as
difficulties in mixing and handling. Below this range, HMPEG
approaches the structure of a conventional nonionic detergent
15 that may impair lather performance and desired user
properties.

 Generally, the polymers will be selected from alkylene
nonionic polymers chemically and terminally attached by
20 hydrophobic moieties, wherein the hydrophobic moiety can be
derivatives of linear or branched alkyl, aryl, alkylaryl,
alkylene, acyl (e.g., having a carbon number of C4 to C40);
fat and oil derivatives of alkylglyceryl, glyceryl, sorbitol,
lanolin oil, coconut oil, jojoba oil, castor oil, almond oil,
25 peanut oil, wheat germ oil, rice bran oil, linseed oil,
apricot pits oil, walnuts, palm nuts, pistachio nuts, sesame
seeds, rapeseed, cade oil, corn oil, peach pit oil, poppyseed
oil, pine oil, soybean oil, avocado oil, sunflower seed oil,
hazelnut oil, olive oil, grapeseed oil, and safflower oil,
30 Shea butter, babassu oil, etc. These hydrophobically
modified polyalkylene glycols are usually commercially
available (see Table 1 for examples).

 To ensure water solubility, it is preferred that the
35 portion of ethylene oxide moiety per mole of HMPEG is between

60% wt. and 99 % wt. (preferably 85 % wt. to 97 % wt.). In other words, the total content of the hydrophobic moiety is between 1 % wt. and 40 % wt. (preferably 3% wt. to 15% wt.) per mole of the defined HMPEG.

5

Specifically, examples of various hydrophobically modified polyalkylene glycols are set forth in Table 1 below where in T_m (°C) were obtained from literature from the corresponding chemical suppliers or measured by the inventors using a differential scanning calorimetry technique.

10

Table 1 Representative hydrophobically modified PEGs.

(R= hydrophobic moieties such as linear or branched alkyl chains (e.g., having carbon number of C4 to C40); derivatives of sorbitol, lanolin radical, coconut radical, jojoba acid radical, castor oil radical, etc.;

POE= Polyoxyethylene (e.g., $-(CH_2CH_2O)_mH$);

m= No. ethylene oxide monomer units; $m>50$.)

10	<u>Chemicals</u>	<u>Suppliers (Brands)</u>	<u>Comments</u>
	POE(m)-R	Witco (Varonic LI-420)	R= glyceryltallowate; m=200; white solid.
15		Seppic (Simusol 220TM)	R=glycerylstearate; m=200; white solid.
		Amerchol (Glucam E-200)	R=glucoside; m=200; water soluble; white solid.
20		Calgene Chemical (600-S)	Tm:52-62C; R=stearate; m=150; Tm:52-62C.
25		Calgene Chemical (600-L)	R=laurate; m=150.
	R-POE(m)-R	Stepan (KESSCO PEG6000 distearate)	R=stearate; m=174; Tm:54C; white solid.
30			

As noted, melting temperature of the compounds is preferred to be about 25°-85°.

35 In addition, the weight ratio of hydrophobically modified polyalkylene glycol to anionic surfactant should be in the range of 1:1.1 to 1:10. While not wishing to be bound by theory, this ratio is believed critical because, at ratios below 1:10, improvement on mildness is not significant and, at ratios above
40 1:1.1, special efforts have to be made to adjust the high viscosity of liquid wash compositions (see the prior art in Patent No. US 4,247,425, US 4,343,726, and US 4,256,611).

The defined hydrophobically modified polyalkylene glycols generally comprise 5 to 25% by wt. (preferably 5 to 15% by wt) of the total liquid composition.

- 5 In addition, the compositions of the invention may include optional ingredients as follows:

 Organic solvents, such as ethanol; auxiliary thickeners, such as carboxymethylcellulose, magnesium aluminum silicate, 10 hydroxyethylcellulose, methylcellulose, carbopols, glucamides, or Antil^(R) from Rhone Poulenc; perfumes; sequestering agents, such as tetrasodium ethylenediaminetetraacetate (EDTA), EHDP or mixtures in an amount of 0.01 to 1%, preferably 0.01 to 0.05%; and coloring agents, opacifiers and pearlizers such as zinc 15 stearate, magnesium stearate, TiO₂, EGMS (ethylene glycol monostearate) or Lytron 621 (Styrene/Acrylate copolymer); all of which are useful in enhancing the appearance or cosmetic properties of the product.

20 The compositions may further comprise antimicrobials such as 2-hydroxy-4,2'4' trichlorodiphenylether (DP300); preservatives such as dimethyloldimethylhydantoin (Glydant XL1000), parabens, sorbic acid etc.

25 The compositions may also comprise coconut acyl mono- or diethanol amides as suds boosters, and strongly ionizing salts such as sodium chloride and sodium sulfate may also be used to advantage.

30 Antioxidants such as, for example, butylated hydroxytoluene (BHT) may be used advantageously in amounts of about 0.01% or higher if appropriate.

35 Cationic conditioners which may be used include Quatrisoft LM-200 Polyquaternium-24, Merquat Plus 3330 - Polyquaternium 39;

and Jaguar^(R) type conditioners.

Polyethylene glycols which may be used include:

5	Polyox	WSR-205	PEG 14M,
	Polyox	WSR-N-60K	PEG 45M, or
	Polyox	WSR-N-750	PEG 7M.

Thickeners which may be used include Amerchol Polymer HM
10 1500 (Nonoxynyl Hydroethyl Cellulose); Glucam DOE 120 (PEG 120 Methyl Glucose Dioleate); Rewoderm^(R) (PEG modified glyceryl cocoate, palmate or tallowate) from Rewo Chemicals; Antil^(R) 141 (from Goldschmidt).

15 Another optional ingredient which may be added are the deflocculating polymers such as are taught in U.S. Patent No. 5,147,576 to Montague, hereby incorporated by reference.

20 Another ingredient which may be included are exfoliants such as polyoxyethylene beads, walnut shells and apricot seeds

The compositions may also contain 0.1 to 15% by wt., preferably 1 to 10% by wt. of a structurant. Such structurants can be used to avoid addition of external structurants (e.g.,
25 cross linked polyacrylates and clays) if suspending particles is desired as well as to provide desirable consumer attributes.

The structurant is generally an unsaturated and/or branched long chain (C₈-C₂₄) liquid fatty acid or ester derivative thereof;
30 and/or unsaturated and/or branched long chain liquid alcohol or ether derivatives thereof. It may also be a short chain saturated fatty acid such as capric acid or caprylic acid. While not wishing to be bound by theory, it is believed that the unsaturated part of the fatty acid or alcohol or the branched
35 part of the fatty acid or alcohol acts to "disorder" the

surfactant hydrophobic chains and induce formation of lamellar phase.

5 Examples of liquid fatty acids which may be used are oleic acid, isostearic acid, linoleic acid, linolenic acid, ricinoleic acid, elaidic acid, arichidonic acid, myristoleic acid and palmitoleic acid. Ester derivatives include propylene glycol isostearate, propylene glycol oleate, glyceryl isostearate, glyceryl oleate and polyglyceryl diisostearate.

10

 Examples of alcohols include oleyl alcohol and isostearyl alcohol. Examples of ether derivatives include isosteareth or oleth carboxylic acid; or isosteareth or oleth alcohol.

15

 The structuring agent may be defined as having melting point below about 25°C centigrade.

 Another optional ingredient is oil/emollient which may be added as a benefit agent to the liquid compositions.

20

 Various classes of oils are set forth below.

25 Vegetable oils: Arachis oil, castor oil, cocoa butter, coconut oil, corn oil, cotton seed oil, olive oil, palm kernel oil, rapeseed oil, safflower seed oil, sesame seed oil and soybean oil.

30 Esters: Butyl myristate, cetyl palmitate, decylolate, glyceryl laurate, glyceryl ricinoleate, glyceryl stearate, glyceryl isostearate, hexyl laurate, isobutyl palmitate, isocetyl stearate, isopropyl isostearate, isopropyl laurate, isopropyl linoleate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, propylene glycol monolaurate, propylene glycol ricinoleate, propylene glycol stearate, and propylene glycol isostearate.

35

Animal Fats: Acetylated lanolin alcohols, lanolin, lard, mink oil and tallow.

5 Fatty acids and alcohols: Behenic acid, palmitic acid, stearic acid, behenyl alcohol, cetyl alcohol, eicosanyl alcohol and isocetyl alcohol.

10 Other examples of oil/emollients include mineral oil, petrolatum, silicone oil such as dimethyl polysiloxane, lauryl and myristyl lactate.

15 It should be understood that where the emollient may also function as a structurant, it should not be doubly included such that, for example, if the structurant is 15% oleyl alcohol, no more than 5% oleyl alcohol as "emollient" would be added since the emollient (whether functioning as emollient or structurant) should not comprise more than 20%, preferably no more than 15% of the composition.

20 The emollient/oil is generally used in an amount from about 1 to 20%, preferably 1 to 15% by wt. of the composition. Generally, it should comprise no more than 20% of the composition.

25 The following examples are intended to illustrate further the invention and are not intended to limit the invention in any way.

30 All percentages are intended to be percentages by weight unless stated otherwise.

EXAMPLES

Protocol

35 Mildness Assessments: Zein dissolution test was used to preliminary screen the irritation potential of the formulations

studied. In an 8 oz. jar, 30 mLs of an aqueous dispersion of a formulation were prepared. The dispersions sat in a 45°C bath until fully dissolved. Upon equilibration at room temperature, 1.5 gms of zein powder were added to each solution with rapid stirring for one hour. The solutions were then transferred to centrifuge tubes and centrifuged for 30 minutes at approximately 3,000 rpms. The undissolved zein was isolated, rinsed and allowed to dry in a 60°C vacuum oven to a constant weight. The percent zein solubilized, which is proportional to irritation potential, was determined gravimetrically.

The Protocol of 3-Day Patch Test: Patch test was used to evaluate skin mildness of aqueous dispersions containing 1% DEFI active (sodium cocoyl isethionate) and different levels of the structurant/coactives. Patches (Hilltop^(R) Chambers, 25 mm in size) were applied to the outer upper arms of the panelists under bandage type dressings (Scanpor^(R) tape). After each designated contact periods (24 hrs. for the first patch application, 18 hrs. for the second and third applications), the patches were removed and the sites were visually ranked in order of severity (erythema and dryness) by trained examiners under consistent lighting.

Formulation Processing: Formulations shows in the examples of this invention were prepared in 400 mL beakers in a 40-60C oil bath. Mixing was accomplished with a variable speed overhead motor. Batch size was varied from 100-250 gms. All chemicals used were commercial materials and used as supplied. Those chemicals were dispersed in Milli-Q water, which accounted for 50-80% of the whole formulation. After the batch was homogeneously mixed, it was allowed to be cooled under room temperature.

Example 1

The irritation reduction potential of hydrophobically modified polyalkylene glycols was investigated using Zein dissolution experiments. As indicated in Figure 1a and Figure 1b, the defined hydrophobically modified polyalkylene glycols (HMPEG), as a class, are significantly more effective than PEG in reducing the Zein % dissolved by 1% to 2.7% aqueous DEFI suspension (DEFI is a sodium acyl isethionate/fatty acid mixture defined in the Table 2 of Example 3). The data in Figure 1a and Figure 1b also showed that at relatively low level of addition of HMPEG (HMPEG to anionic surfactant weight ratio below 1:1), HMPEG significantly reduced the amount of Zein dissolved by DEFI.

Example 2

Three day skin patch tests showed that, even at low levels of addition, POE(200) glyceryl stearate significantly reduced the skin irritation caused by even a mild surfactant system, such as sodium acyl isethionate/cocoamido propyl betaine. As shown in Figure 2, at a POE(200) glyceryl stearate to sodium acyl isethionate (SAI) weight ratio around 0.74:1 (equivalent to 10% wt. POE(200) glyceryl stearate in a liquid containing 13.5% wt. sodium acyl isethionate), POE(200) glyceryl stearate reduced the skin irritation of a DEFI/betaine liquor significantly. In contrast, even at a PEG 8000 to SAI weight ratio as high as 1.67:1 (effectively 22.5% wt. PEG 8000 in a liquid with 13.5% SAI), PEG 8000 made no measurable mildness contribution to the SAI/CAP betaine aqueous liquor.

Example 3

All amounts are given in percentage of weight. These formulations used sodium cocoyl isethionate as the major anionic detergent with other amphoteric and anionic surfactants as

coactives. The formulation (A) was a stable milky white cream, which provided rich, creamy, and slippery lather that was rinsed off easily. The formulation (B) and (C) were stable milky white lotions that were pourable and pumpable. These lotions provided rich and creamy lather.

Table 2

Formulation	(A)	(B)	(C)
Sodium cocoyl isethionate (From DEFI*)	0	0	14.5%
Sodium cocoyl isethionate (From IGEPON AC-78)	10.0%	9.0%	0.0
Cocoamidopropyl betaine	5.0	4.5	3.8
Sodium lauryl ether sulphate, 3EO	0.0	1.8	4.8
Glycerin	0.0	1.4	1.0
Palmitic-stearate acid (From IGEPON or DEFI)	0.4	0.4	4.5
POE(200) glyceryl stearate	5.0	0.0	6.0
POE(200) glyceryl tallowate	0.0	7.0	0.0
Propylene glycol	0.0	4.8	0.0
Sodium chloride	2.0	1.8	1.4
Ammonium chloride	0.0	5.8	5.0
Sodium isethionate	0.4	0.4	0.2
Water	balance to 100%	balance to 100%	balance to 100%

*DEFI: directly esterified fatty acid isethionate, which is a mixture containing about 74% by weight of sodium acyl isethionate, 23% stearic-palmitic acid and small amounts of other materials, manufactured by Lever Brothers Co, U.S.

Example 4

All amounts are given in percentage of weight. These formulations used sodium lauryl sulphate. (3EO) as the major

anionic detergent with optional amphoteric and anionic surfactants as coactives. These clear, pourable liquids provided rich, creamy and slippery lather and smooth skin feel.

5

Table 3

10

15

Formulation	(C)	(D)	(E)
SLES (3EO)	5.0	10.0	15.0
Sodium lauryl sarcosinate	5.0	3.0	0.0
Cocoamidopropyl betaine	5.0	5.0	10.0
Propylene glycol	2.0	1.0	2.0
POE(200) glyceryl stearate	5.0	10.0	10.0
Water	Balance to 100%	Balance to 100%	Balance to 100%

CLAIMS

1. A liquid detergent composition comprising:

(a) a detergent surfactant system comprising

5 (1) 3% to 30% by wt. total composition anionic or mixtures of anionic surfactants wherein the anionic surfactant comprises 50% or greater of the detergent active system; and

10 (2) 0.1% to 20% by wt. total composition comprising one or more amphoteric surfactants, and

(b) 5 to 25% by wt. hydrophobically modified polyalkylene glycols;

15 wherein the ratio of hydrophobically modified polyalkylene glycol to anionic or anionics is 1:1.1 to 1:10; and

20 wherein the hydrophobically modified polyalkylene glycol is defined by having a molecular weight of 4,000 to 25,000 Dalton; having a weight percentage of alkylene oxide comprising the polymer between 60% and 99%; and having a melting temperature of 25 °C to 85 °C.

25 2. A composition as claimed in claim 1, wherein said anionic surfactant or surfactants comprises 5% to 20% of the composition.

3. A composition as claimed in claim 1 or 2, wherein said amphoteric surfactant comprises 3% to 10% of the composition.

30 4. A composition as claimed in any preceding claim, wherein weight ratio of hydrophobically modified polyalkylene glycol to anionic surfactants is between 1:1.1 to 1:7.

35 5. A composition as claimed in any preceding claim, wherein alkylene oxide comprises 85% to 97% by wt. of said hydrophobically modified polyalkylene glycol.

6. A composition as claimed in any preceding claim, wherein MW of said hydrophobically modified polyalkylene glycol is between 4,000 to 15,000 Dalton.

- 5 7. A method of enhancing mildness of a liquid detergent composition comprising a surfactant system comprising 3% to 30% by wt. anionic or mixture of anionics which method comprises adding 5 to 25% by wt. of an hydrophobically modified polyalkylene glycol to said composition;

10

wherein the ratio of hydrophobically modified polyalkylene glycol to anionic or anionics is 1:1.1 to 1:10;

- 15 wherein the hydrophobically modified polyalkylene glycol is defined by having the percentage of alkylene oxide comprising the hydrophobically modified polyalkylene glycol between 60% and 99%; and as having a molecular weight of 4,000 to 25,000 Dalton.

Figure 1 Dose response of POE(200)-glyceryl-Stearate/-tallowate to the Zein% dissolved in DEFI/CAP betaine aqueous liquors

Figure 1a

1.08% DEFI (0.8% SAI), 0.2% CAP Betaine




Figure 1b

2.7% DEFI (2% SAI), 0.27% CAP Betaine





Figure 2 Patch Tests: Effect of Hydrophobically Modified PEG on DEFI/Cap Betaine [29]

Effect of HMPEG on irritation potential of an anionic/amphoteric surfactant system
(1% wt. sodium acyl rsethionate (SAI) / 0.26% wt. Cocoamido propyl betaine (control))



INTERNATIONAL SEARCH REPORT

 Inv. Application No
 PCT/EP 97/01147

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C11D1/94 C11D3/37 A61K7/50 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 04882 A (PROCTER & GAMBLE) 2 April 1992 see page 10, line 6 - page 11, line 72; claims 9,10; examples 4-8 ---	1-7
X	WO 93 15711 A (PROCTER & GAMBLE) 19 August 1993 see page 10, line 9 - page 11, line 12; claims; examples 1-5 ---	1-7
A	US 4 426 310 A (VERUNICA PIERRE M) 17 January 1984 see claim 5; examples ---	1-7
A	US 4 247 425 A (EGAN RICHARD R ET AL) 27 January 1981 cited in the application see the whole document ---	1,2,7
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

4 September 1997

Date of mailing of the international search report

15.09.97

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Grittern, A

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 97/01147

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 450 091 A (SCHMOLKA IRVING R) 22 May 1984 see claims; examples 1-7 ---	1,2,4
A	JP 54 043 212 A (LION CORP) 5 April 1979 see page 70; table 3 ---	1-3,5,6
A	JP 04 122 797 A (SHISEIDO CO LTD) 23 April 1992 see page 704, line 15 - line 25; tables -----	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 97/01147

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9204882 A	02-04-92	AU 8730291 A	15-04-92
		CN 1060782 A	06-05-92
		CZ 9300453 A	16-02-94
		DE 69115282 D	18-01-96
		DE 69115282 T	18-07-96
		EP 0548265 A	30-06-93
		ES 2080962 T	16-02-96
		HU 64206 A	28-12-93
		JP 6501011 T	27-01-94
		SK 22193 A	07-07-93
		TR 26415 A	15-03-95
WO 9315711 A	19-08-93	AU 3598493 A	03-09-93
		CA 2129129 A	19-08-93
		CN 1078631 A	24-11-93
		EP 0625040 A	23-11-94
		JP 7503731 T	20-04-95
US 4426310 A	17-01-84	AU 554574 B	28-08-86
		AU 8165682 A	15-09-83
US 4247425 A	27-01-81	US 4343726 A	10-08-82
US 4450091 A	22-05-84	CA 1210701 A	02-09-86
JP 54043212 A	05-04-79	JP 60009693 B	12-03-85
JP 04122797 A	23-04-92	NONE	